



Fairfax Oral and  
Maxillofacial Surgery

Patrick J. Dolan D.D.S., M.S.

Kimberly A. Silloway, D.D.S.

Jonathan J. Park, D.M.D., M.D.

Daniel M. Theberge, D.D.S.

Cyrus A. Ramsey, D.M.D., M.D.

Steven F. Robertson, D.D.S.

Frederick Liu, D.D.S., M.D.

Justin A. Baker, D.D.S.

John H. Hur, D.D.S., M.D.

Henry W. Zussman, D.D.S.\*

Peter W. Smith, D.D.S.\*

Jeffrey A. Lane, D.D.S.\*

Diplomates of the American Board  
of Oral and Maxillofacial Surgery

**Fairfax**  
10530 Rosehaven Street, Suite 111  
Fairfax, VA 22030  
(703) 385-5777

**Centreville**  
6211 Centreville Road, Suite 800  
Centreville, VA 20121  
(703) 266-0026

**Reston**  
1800 Towne Center Drive, Suite 416  
Reston, VA 20190  
(703) 435-7100

**Burke**  
5252 Lyngate Court, Suite 101  
Burke, VA 22015  
(703) 764-2077

**Woodbridge**  
3421 Commission Court, Suite 202  
Woodbridge, VA 22192  
(703) 672-4200

[www.fairfaxoralsurgery.com](http://www.fairfaxoralsurgery.com)

## DISCHARGE INSTRUCTIONS FOR BONE GRAFT PROCEDURES

The *bone graft* procedure entails surgically placing bone or bone material.

### What to expect after surgery:

- *Swelling and bruising (May take 24-48 hours to peak)*
- *Bleeding*
- *Numbness due to anesthesia for up to six hours*
- *Discomfort for several days*
- *Facial and sinus pressure*
- *You may notice white particles within the mouth several days after surgery*
- *If suture was placed, it will dissolve*
- *If screws were placed, these will be removed at a following appointment, by discretion of the doctor*

### Please follow these instructions:

- *Drink plenty of liquids*
- *Bleeding – lasts several hours after surgery – place gauze in area for 20-30 minutes. Repeat as needed.*
- *Avoid salty, spicy, crunchy and citric foods*
- *Avoid mouthwashes – rinse with saltwater three times per day for one week.*
- *Avoid hot foods or liquids while still numb*
- *To reduce swelling, elevate head and apply ice to the face, use moist heat to reduce swelling after the first 24 hours*
- *Eat a soft diet for the first 24 hours, avoid chewing in the surgical site*
- *Soft diet examples- yogurt, pancakes, smoothies, eggs, mashed potatoes*

### Antibiotics and Pain Medications

Have any prescriptions filled promptly and take as directed. Non-narcotic medications (*Motrin, Advil, Ibuprofen, Tylenol*) should be taken as soon as possible and repeated as prescribed for the first several days. Narcotic pain medication can be taken in addition if pain relief is not adequate. However, when narcotic pain medications (*Percocet, Vicodin, Tylenol with Codeine*) are taken, you should not operate automobiles or other power equipment. Antibiotics, when prescribed, should be taken as indicated until gone.

If a rash or swelling develops, discontinue the antibiotic immediately and call the office to inform your doctor.

### Oral Hygiene and Smoking

It is important to maintain oral hygiene so that the wound will heal properly and not become infected. Smokers are advised not to smoke for five to seven days after surgery. Drawing on a cigarette may disturb the blood clot and thus interfere with the healing process.

### Physical Activities

Please reduce activities for 3 days following surgery.

### Call your Doctor if:

- *You have signs of infection, sudden swelling, or fever*
- *You have a reaction to any of the medications*
- *You experience prolonged bleeding*
- *Pain increases and/or your condition worsens*
- *You have any questions regarding the surgery or post-surgery instructions. If calling after business hours, you will be directed to an after-hours answering service at 855-808-9500 and a doctor will promptly return your call*

\*Emeritus

Discharged by: