



Fairfax Oral and
Maxillofacial Surgery

Patrick J. Dolan D.D.S., M.S.

Kimberly A. Silloway, D.D.S.

Jonathan J. Park, D.M.D., M.D.

Daniel M. Theberge, D.D.S.

Cyrus A. Ramsey, D.M.D., M.D.

Steven F. Robertson, D.D.S.

Frederick Liu, D.D.S., M.D.

Justin A. Baker, D.D.S.

John H. Hur, D.D.S., M.D.

Henry W. Zussman, D.D.S.*

Peter W. Smith, D.D.S.*

Jeffrey A. Lane, D.D.S.*

Diplomates of the American Board
of Oral and Maxillofacial Surgery

Fairfax
10530 Rosehaven Street, Suite 111
Fairfax, VA 22030
(703) 385-5777

Centreville
6211 Centreville Road, Suite 800
Centreville, VA 20121
(703) 266-0026

Reston
1800 Towne Center Drive, Suite 416
Reston, VA 20190
(703) 435-7100

Burke
5252 Lyngate Court, Suite 101
Burke, VA 22015
(703) 764-2077

Woodbridge
3421 Commission Court, Suite 202
Woodbridge, VA 22192
(703) 672-4200

www.fairfaxoralsurgery.com

*Emeritus

DISCHARGE INSTRUCTIONS FOR EXPOSE AND BOND PROCEDURE

The *expose and bond* procedure entails exposing an impacted tooth and bonding an orthodontic bracket to assist in your orthodontic treatment.

What to expect after surgery:

- *Slight swelling*
- *Minor bleeding*
- *Numbness due to anesthesia for up to six hours*
- *Discomfort for several days*
- *Sutures will be placed and will dissolve*
- *A gold or elastic chain will be visible from the surgical site and will attach to your existing braces*

Please follow these instructions:

- *Drink plenty of liquids*
- *Eat a soft diet for the first 24 hours*
- *Avoid hot foods or liquids while still numb*
- *Avoid crunchy foods for the first week*
- *If bleeding occurs, apply firm pressure to the area for 15-20 minutes with cotton rolls or gauze that was provided. Repeat as needed.*
- *Avoid mouthwashes – rinse with saltwater three times per day for one week.*
- *To reduce swelling, elevate head and apply ice packs to the face*
- *Limit physical activity the day of surgery.*

Antibiotics and Pain Medications

Have any prescriptions filled promptly and take as directed. Non-narcotic medications (*Motrin, Advil, Ibuprofen, Tylenol*) should be taken as soon as possible and repeated as prescribed for the first several days.

Narcotic pain medication can be taken in addition if pain relief is not adequate. However, when narcotic pain medications (*Percocet, Vicodin, Tylenol with Codeine*) are taken, you should not operate automobiles or other power equipment. Antibiotics, when prescribed, should be taken as indicated until gone. If a rash or swelling develops, discontinue the antibiotic immediately and call the office to inform your doctor.

Oral Hygiene and Smoking

It is important to maintain oral hygiene so that the wound will heal properly and not become infected.

Call your Doctor if:

- *The gold or elastic chain or bracket loosens and comes off.*
- *You have signs of infection, sudden swelling, or fever*
- *You have a reaction to any of the medications*
- *You experience prolonged bleeding*
- *Pain increases and/or your condition worsens*
- *You have any questions regarding the surgery or post-surgery instructions. If calling after business hours, you will be directed to an after-hours answering service at 855-808-9500 and a doctor will promptly return your call*

Discharged by:
